

The Many Phases of Massive Galaxies

A Near-Infrared Spectroscopic Study of Galaxies in the Early Universe

The Many Phases of Massive Galaxies

A Near-Infrared Spectroscopic Study of Galaxies in the Early Universe

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. P. F. van der Heijden,
volgens besluit van het College voor Promoties
te verdedigen op woensdag 26 september 2007
klokke 16.15 uur

door

Mariska Therese Kriek

geboren te Leiden
in 1979

Promotiecommissie

Promotores: Prof. dr. M. Franx
Prof. dr. P. G. van Dokkum (Yale University, USA)

Referent: Prof. dr. P. Barthel (Rijksuniversiteit Groningen)

Overige leden: Prof. dr. P. T. de Zeeuw
Prof. dr. G. D. Illingworth (University of California, Santa Cruz, USA)
Prof. dr. K. Kuijken
Dr. P. P. van der Werf

Table of contents

	Page
1 Introduction	1
1.1 Structure Formation	1
1.2 Galaxy Properties	1
1.3 Massive Galaxies over Cosmic Time	2
1.4 This Thesis	4
1.5 Conclusions and Outlook	6
2 Stellar Continua and Balmer/4000Å Breaks of Red $z > 2$ Galaxies	9
2.1 Introduction	10
2.2 Data	11
2.2.1 Target Selection and Photometry	11
2.2.2 Observations	12
2.2.3 Reduction of GNIRS Spectra	13
2.2.4 Extraction of One-dimensional Spectra	14
2.3 Spectral Modeling	16
2.3.1 Fitting Procedure and Results	17
2.3.2 Continuum Redshifts	19
2.3.3 Comparison to Broadband SEDs	20
2.3.4 Discussion of Modeling Results	20
2.4 Spectral Diagnostics	21
2.4.1 Break Indices	21
2.4.2 Comparison to $H\alpha$ Equivalent Width	22
2.4.3 Active Galactic Nuclei?	23
2.5 Comparison to Other Galaxies	24
2.5.1 Comparison to Other High-Redshift Galaxies	24
2.5.2 Comparison with Low-Redshift Galaxies	25
2.6 Summary and Conclusions	26
3 Massive Galaxies at $z \sim 2.3$ with Strongly Suppressed Star Formation	29
3.1 Introduction	30
3.2 Galaxy Sample and Data	30
3.3 Suppressed Star Formation	32
3.4 Discussion	34
4 The Origin of Line Emission in Massive $z \sim 2.3$ Galaxies	37
4.1 Introduction	38

4.2	Data	39
4.2.1	Sample	39
4.2.2	GNIRS Spectra	40
4.2.3	SINFONI Spectra	41
4.2.4	Extraction of One-Dimensional Spectra	42
4.2.5	Line Measurements	44
4.2.6	Line Maps	46
4.3	Origin of the Line Emission	47
4.3.1	Emission Line Ratios	47
4.3.2	Other Diagnostics of AGN Activity	49
4.3.3	Summary and AGN Fraction	52
4.4	Implications	53
4.4.1	Stellar Populations of AGN Host Galaxies	53
4.4.2	Stellar Masses of AGN Host Galaxies	55
4.4.3	Downsizing of AGN Host Galaxies	56
4.4.4	AGNs and the Suppression of Star Formation	58
4.5	Summary and Conclusions	59
5	A Near-Infrared Spectroscopic Survey of K-selected Galaxies at $z \sim 2.3$	65
5.1	Introduction	66
5.2	Data	67
5.2.1	Sample Selection	67
5.2.2	NIR spectra	70
5.3	Spectroscopic versus Photometric Redshifts	75
5.3.1	Spectroscopic Redshifts and Galaxy Properties	75
5.3.2	Photometric Redshifts and Galaxy Properties	79
5.3.3	Direct Comparison	79
5.3.4	Systematics and Catastrophic Failures	80
5.3.5	An Empirically-Motivated Template Set	83
5.4	Implications	84
5.4.1	Implications for Derived Properties	84
5.4.2	Implications for Photometric Samples	87
5.4.3	Implications for Previous Studies	91
5.5	Summary	92
6	The Detection of a Red Sequence at $z \sim 2.3$ and its Evolution to $z \sim 0$	97
6.1	Introduction	98
6.2	Data	99
6.3	A Red Sequence at $z \sim 2.3$	99
6.3.1	The Detection of the Red Sequence at $z \sim 2.3$	99
6.3.2	Properties of Red-Sequence Galaxies	101
6.4	Measuring the Evolution of the Red Sequence	104
6.4.1	Spectroscopic Samples at Lower Redshifts	104
6.4.2	The Color Evolution of the Red Sequence	104
6.4.3	Evolution of the Mass and Number Density	108

6.5	Modeling the Evolution of the Red Sequence	109
6.5.1	Aging of Stellar populations	109
6.5.2	Growth of the Red Sequence	109
6.5.3	Red Mergers	111
6.5.4	Other Influences	112
6.6	Summary and Conclusions	112
	Nederlandse Samenvatting	115
	Curriculum Vitae	121
	Nawoord	123

