

## Bibliography

- Al-Adhami MA K. Y. (1977) Ontogenesis of haematopoietic sites in *Brachydanio rerio* (hamilton-buchanan) teleostei. *Dev Growth Differ* **19**, 171-9.
- Arnold I. and Watt F. M. (2001) c-Myc activation in transgenic mouse epidermis results in mobilization of stem cells and differentiation of their progeny. *Curr Biol* **11**, 558-68.
- Barclay A. N. (2003) Membrane proteins with immunoglobulin-like domains--a master superfamily of interaction molecules. *Semin Immunol* **15**, 215-23.
- Barreto V. M., Pan-Hammarstrom Q., Zhao Y., Hammarstrom L., Misulovin Z. and Nussenzweig M. C. (2005) AID from bony fish catalyzes class switch recombination. *J Exp Med* **202**, 733-8.
- Berman J., Hsu K. and Look A. T. (2003) Zebrafish as a model organism for blood diseases. *Br J Haematol* **123**, 568-76.
- Blanpain C., Lowry W. E., Pasolli H. A. and Fuchs E. (2006) Canonical notch signaling functions as a commitment switch in the epidermal lineage. *Genes Dev* **20**, 3022-35.
- Blom N., Gammeltoft S. and Brunak S. (1999) Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *J Mol Biol* **294**, 1351-62.
- Buljan M. and Bateman A. (2009) The evolution of protein domain families. *Biochem Soc Trans* **37**, 751-5.
- Bushell K. M., Sollner C., Schuster-Boeckler B., Bateman A. and Wright G. J. (2008) Large-scale screening for novel low-affinity extracellular protein interactions. *Genome Res* **18**, 622-30.
- Cagan R. L. and Ready D. F. (1989) Notch is required for successive cell decisions in the developing *Drosophila* retina. *Genes Dev* **3**, 1099-112.
- Cao T., Racz P., Szauder K. M., Groma G., Nakamatsu G. Y., Fogelgren B., Pankotai E., He Q. P. and Csizsar K. (2007) Mutation in *MpZ3*, a gene encoding a predicted the adhesion protein, in the rough coat (*rc*) mice with severe skin and hair abnormalities. *J Invest Dermatol* **127**, 1375-86.
- Carenini S., Maurer M., Werner A., Blazyca H., Toyka K. V., Schmid C. D., Raivich G. and Martini R. (2001) The role of macrophages in demyelinating peripheral nervous system of mice heterozygously deficient in *po*. *J Cell Biol* **152**, 301-8.
- Chapon F., Latour P., Diraison P., Schaeffer S. and Vandenberghe A. (1999) Axonal phenotype of Charcot-Marie-Tooth disease associated with a mutation in the myelin protein zero gene. *J Neurol Neurosurg Psychiatry* **66**, 779-82.
- Chow R. L., Altmann C. R., Lang R. A. and Hemmati-Brivanlou A. (1999) *Pax6* induces ectopic eyes in a vertebrate. *Development* **126**, 4213-22.
- Corripio-Miyar Y., Bird S., Treasurer J. W. and Secombes C. J. (2007) *RAG-1* and *IgM* genes, markers for early development of the immune system in the gadoid haddock, *Melanogrammus aeglefinus*, L. *Fish Shellfish Immunol* **23**, 71-85.
- Crispin J. C. and Tsokos G. C. (2008) Novel molecular targets in the treatment of systemic lupus erythematosus. *Autoimmun Rev* **7**, 256-61.
- Czerny T., Halder G., Kloter U., Souabni A., Gehring W. J. and Busslinger M. (1999) twin of *eyeless*, a second *Pax-6* gene of *Drosophila*, acts upstream of *eyeless* in the control of eye development. *Mol Cell* **3**, 297-307.
- Davies D. R., Sarma R., Labaw L. W., Silvertown E., Segal D. and Terry W. D. (1971) X-ray diffraction and electron microscope studies on a crystalline human immunoglobulin. *Ann N Y Acad Sci* **190**, 122-9.
- De Jonghe P., Timmerman V., Ceuterick C., Nelis E., De Vriendt E., Lofgren A., Vercruyssen A., Verellen C., Van Maldergem L., Martin J. J. and Van Broeckhoven C. (1999) The Thr124Met mutation in the peripheral myelin protein zero (*MPZ*) gene is associated with a clinically distinct Charcot-Marie-Tooth phenotype. *Brain* **122** (Pt 2), 281-90.
- Demehri S. and Kopan R. (2009) Notch signaling in bulge stem cells is not required for selection of hair follicle fate. *Development* **136**, 891-6.
- DeMonte L., Porcellini S., Tafi E., Sheridan J., Gordon J., DePreter M., Blair N., Panigada M., Sanvito F., Merati B., Albientz A., Barthlott T., Ozmen L., Blackburn C. C. and Guttinger M. (2007) *EVA* regulates thymic stromal organisation and early thymocyte development. *Biochem Biophys Res Commun* **356**, 334-40.
- Dickie M. (1966) Rough coat. *Mouse News Letter* **34**.
- Eicher EM F., S., Reynolds, S. (1977) Rough coat on Chromosome 9. *Mouse News Letter* **56**.
- Fernandez-Gonzalez A., La Spada A. R., Treadaway J., Higdon J. C., Harris B. S., Sidman R. L., Morgan J. I. and Zuo J. (2002) Purkinje cell degeneration (*pcd*) phenotypes caused by mutations in the axotomy-induced gene, *Nnai*. *Science* **295**, 1904-6.
- Fraser J. S., Yu Z., Maxwell K. L. and Davidson A. R. (2006) Ig-like domains on bacteriophages: a tale of promiscuity and deceit. *J Mol Biol* **359**, 496-507.
- Freytschmidt-Paul P., Hoffmann R., Levine E., Sundberg J. P., Happle R. and McElwee K. J. (2001) Current and potential agents for the treatment of alopecia areata. *Curr Pharm Des* **7**, 213-30.
- Frye M., Gardner C., Li E. R., Arnold I. and Watt F. M. (2003) Evidence that Myc activation depletes the epidermal stem cell compartment by modulating adhesive interactions with the local microenvironment. *Development* **130**, 2793-808.

- Gage P. J. and Zacharias A. L. (2009) Signaling “cross-talk” is integrated by transcription factors in the development of the anterior segment in the eye. *Dev Dyn* **238**, 2149-62.
- Gaiano N. and Fishell G. (2002) The role of notch in promoting glial and neural stem cell fates. *Annu Rev Neurosci* **25**, 471-90.
- Gilhar A. and Kalish R. S. (2006) Alopecia areata: a tissue specific autoimmune disease of the hair follicle. *Autoimmun Rev* **5**, 64-9.
- Gilhar A., Kam Y., Assy B. and Kalish R. S. (2005) Alopecia areata induced in C<sub>3</sub>H/HeJ mice by interferon-gamma: evidence for loss of immune privilege. *J Invest Dermatol* **124**, 288-9.
- Ginalski K., Elofsson A., Fischer D. and Rychlewski L. (2003) 3D-Jury: a simple approach to improve protein structure predictions. *Bioinformatics* **19**, 1015-8.
- Gompel N., Cubedo N., Thisse C., Thisse B., Dambly-Chaudiere C. and Ghysen A. (2001) Pattern formation in the lateral line of zebrafish. *Mech Dev* **105**, 69-77.
- Grillitsch S., Medgyesy N., Schwerte T. and Pelster B. (2005) The influence of environmental P(O<sub>2</sub>) on hemoglobin oxygen saturation in developing zebrafish *Danio rerio*. *J Exp Biol* **208**, 309-16.
- Guex N. a. P., M.C. (1997) SM2 SWISS-MODEL and the Swiss-PdbViewer: An environment for comparative protein modeling. *Electrophoresis* **18**, 2714-2723.
- Gupta R. J. E., Brunak S. (2004) Prediction of N-glycosylation sites in human proteins. *In preparation*.
- Guttinger M., Sutti F., Panigada M., Porcellini S., Merati B., Mariani M., Teesalu T., Consalez G. G. and Grassi F. (1998) Epithelial V-like antigen (EVA), a novel member of the immunoglobulin superfamily, expressed in embryonic epithelia with a potential role as homotypic adhesion molecule in thymus histogenesis. *J Cell Biol* **141**, 1061-71.
- Halaby D. M. and Mornon J. P. (1998) The immunoglobulin superfamily: an insight on its tissular, species, and functional diversity. *J Mol Evol* **46**, 389-400.
- Harpaz Y. and Chothia C. (1994) Many of the immunoglobulin superfamily domains in cell adhesion molecules and surface receptors belong to a new structural set which is close to that containing variable domains. *J Mol Biol* **238**, 528-39.
- Hawkins R. D., Bashiardes S., Powder K. E., Sajan S. A., Bhonagiri V., Alvarado D. M., Speck J., Warchol M. E. and Lovett M. (2007) Large scale gene expression profiles of regenerating inner ear sensory epithelia. *PLoS One* **2**, e525.
- Hayashi K., Cao T., Passmore H., Jourdan-Le Saux C., Fogelgren B., Khan S., Hornstra I., Kim Y., Hayashi M. and Csiszar K. (2004) Progressive hair loss and myocardial degeneration in rough coat mice: reduced lysyl oxidase-like (LOXL) in the skin and heart. *J Invest Dermatol* **123**, 864-71.
- Horiguchi M., Inoue T., Ohbayashi T., Hirai M., Noda K., Marmorstein L. Y., Yabe D., Takagi K., Akama T. O., Kita T., Kimura T. and Nakamura T. (2009) Fibulin-4 conducts proper elastogenesis via interaction with cross-linking enzyme lysyl oxidase. *Proc Natl Acad Sci U S A* **106**, 19029-34.
- Hutchinson E. G. and Thornton J. M. (1993) The Greek key motif: extraction, classification and analysis. *Protein Eng* **6**, 233-45.
- Jiang Y. J., Brand M., Heisenberg C. P., Beuchle D., Furutani-Seiki M., Kelsh R. N., Warga R. M., Granato M., Haffter P., Hammerschmidt M., Kane D. A., Mullins M. C., Odenthal J., van Eeden F. J. and Nusslein-Volhard C. (1996) Mutations affecting neurogenesis and brain morphology in the zebrafish, *Danio rerio*. *Development* **123**, 205-16.
- Julenius K., Molgaard A., Gupta R. and Brunak S. (2005) Prediction, conservation analysis, and structural characterization of mammalian mucin-type O-glycosylation sites. *Glycobiology* **15**, 153-64.
- Katoh M. (2008) WNT signaling in stem cell biology and regenerative medicine. *Curr Drug Targets* **9**, 565-70.
- Kazakova N., Li H., Mora A., Jessen K. R., Mirsky R., Richardson W. D. and Smith H. K. (2006) A screen for mutations in zebrafish that affect myelin gene expression in Schwann cells and oligodendrocytes. *Dev Biol* **297**, 1-13.
- Kerstetter A. E., Azodi E., Marrs J. A. and Liu Q. (2004) Cadherin-2 function in the cranial ganglia and lateral line system of developing zebrafish. *Dev Dyn* **230**, 137-43.
- Korhonen J., Partanen J., Armstrong E., Vaahtokari A., Elenius K., Jalkanen M. and Alitalo K. (1992) Enhanced expression of the tie receptor tyrosine kinase in endothelial cells during neovascularization. *Blood* **80**, 2548-55.
- Krogh A., Larsson B., von Heijne G. and Sonnhammer E. L. (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**, 567-80.
- Kuchler A. M., Gjini E., Peterson-Maduro J., Cancilla B., Wolburg H. and Schulte-Merker S. (2006) Development of the zebrafish lymphatic system requires VEGFC signaling. *Curr Biol* **16**, 1244-8.
- Lathia J. D., Mattson M. P. and Cheng A. (2008) Notch: from neural development to neurological disorders. *J Neurochem* **107**, 1471-81.
- Lee J., Basak J. M., Demehri S. and Kopan R. (2007) Bi-compartmental communication contributes to the opposite proliferative behavior of Notch1-deficient hair follicle and epidermal keratinocytes. *Development* **134**, 2795-806.
- Letunic I., Goodstadt L., Dickens N. J., Doerks T., Schultz J., Mott R., Ciccarelli F., Copley R. R., Ponting C. P. and Bork P. (2002) Recent improvements to the SMART domain-based sequence annotation resource. *Nucleic Acids Res* **30**, 242-4.

- Leyns L., Bouwmeester T., Kim S. H., Piccolo S. and De Robertis E. M. (1997) Frzb-1 is a secreted antagonist of Wnt signaling expressed in the Spemann organizer. *Cell* **88**, 747-56.
- Loosli F., Winkler S. and Wittbrodt J. (1999) Six3 overexpression initiates the formation of ectopic retina. *Genes Dev* **13**, 649-54.
- Ma A., Koka R. and Burkett P. (2006) Diverse functions of IL-2, IL-15, and IL-7 in lymphoid homeostasis. *Annu Rev Immunol* **24**, 657-79.
- Ma E. Y. and Raible D. W. (2009) Signaling pathways regulating zebrafish lateral line development. *Curr Biol* **19**, R381-6.
- Manly K. F., Cudmore R. H., Jr. and Meer J. M. (2001) Map Manager QTX, cross-platform software for genetic mapping. *Mamm Genome* **12**, 930-2.
- Martinez-Mir A., Zlotogorski A., Gordon D., Petukhova L., Mo J., Gilliam T. C., Londono D., Haynes C., Ott J., Hordinsky M., Nanova K., Norris D., Price V., Duvic M. and Christiano A. M. (2007) Genomewide scan for linkage reveals evidence of several susceptibility loci for alopecia areata. *Am J Hum Genet* **80**, 316-28.
- Maurer M., Kobsar I., Berghoff M., Schmid C. D., Carenini S. and Martini R. (2002) Role of immune cells in animal models for inherited neuropathies: facts and visions. *J Anat* **200**, 405-14.
- McDonagh A. J. and Tazi-Ahni R. (2002) Epidemiology and genetics of alopecia areata. *Clin Exp Dermatol* **27**, 405-9.
- McMillan J. R. and Shimizu H. (2001) Desmosomes: structure and function in normal and diseased epidermis. *J Dermatol* **28**, 291-8.
- Meeker N. D. and Trede N. S. (2008) Immunology and zebrafish: spawning new models of human disease. *Dev Comp Immunol* **32**, 745-57.
- Metaserver1. <http://bioinfo.pl/meta/>.
- Miller J. R. (2002) The Wnts. *Genome Biol* **3**, REVIEWS3001.
- Misu K., Yoshihara T., Shikama Y., Awaki E., Yamamoto M., Hattori N., Hirayama M., Takegami T., Nakashima K. and Sobue G. (2000) An axonal form of Charcot-Marie-Tooth disease showing distinctive features in association with mutations in the peripheral myelin protein zero gene (Thr124Met or Asp75Val). *J Neurol Neurosurg Psychiatry* **69**, 806-11.
- Moog-Lutz C., Cave-Riant F., Guibal F. C., Breaux M. A., Di Gioia Y., Couraud P. O., Cayre Y. E., Bourdoulous S. and Lutz P. G. (2003) JAML, a novel protein with characteristics of a junctional adhesion molecule, is induced during differentiation of myeloid leukemia cells. *Blood* **102**, 3371-8.
- Mulroy T., McMahan J. A., Burakoff S. J., McMahan A. P. and Sen J. (2002) Wnt-1 and Wnt-4 regulate thymic cellularity. *Eur J Immunol* **32**, 967-71.
- Niehrs C. (1999) Head in the WNT: the molecular nature of Spemann's head organizer. *Trends Genet* **15**, 314-9.
- Park Y. G., Hayasaka S., Takagishi Y., Inouye M., Okumoto M. and Oda S. (2001) Histological characteristics of the pelage skin of rough fur mice (C3H/HeJ- ruf/ruf). *Exp Anim* **50**, 179-82.
- Pelster B. and Burggren W. W. (1996) Disruption of hemoglobin oxygen transport does not impact oxygen-dependent physiological processes in developing embryos of zebra fish (*Danio rerio*). *Circ Res* **79**, 358-62.
- Peri S. e. a. (2003) Development of human protein reference database as an initial platform for approaching systems biology in humans. *Genome Research* **13**, 2363-2371.
- Poljak R. J., Amzel L. M., Avey H. P., Chen B. L., Phizackerley R. P. and Saul F. (1973) Three-dimensional structure of the Fab' fragment of a human immunoglobulin at 2,8-Å resolution. *Proc Natl Acad Sci U S A* **70**, 3305-10.
- Racz P., Mink M., Ordas A., Cao T., Szalma S., Szauder K. M. and Csiszar K. (2009) The human orthologue of murine Mpz3 with predicted adhesive and immune functions is a potential candidate gene for immune-related hereditary hair loss. *Exp Dermatol* **18**, 261-3.
- Rebagliati M. R., Toyama R., Haffter P. and Dawid I. B. (1998) cyclops encodes a nodal-related factor involved in midline signaling. *Proc Natl Acad Sci U S A* **95**, 9932-7.
- Richardson J. S. (1981) The anatomy and taxonomy of protein structure. *Adv Protein Chem* **34**, 167-339.
- Ruvinsky I., Chertkov O., Borue X. V., Agulnik S. I., Gibson-Brown J. J., Lyle S. R. and Silver L. M. (2002) Genetics analysis of mouse mutations Abnormal feet and tail and rough coat, which cause developmental abnormalities and alopecia. *Mamm Genome* **13**, 675-9.
- Schmid C. D., Stienekemeier M., Oehen S., Bootz F., Zielasek J., Gold R., Toyka K. V., Schachner M. and Martini R. (2000) Immune deficiency in mouse models for inherited peripheral neuropathies leads to improved myelin maintenance. *J Neurosci* **20**, 729-35.
- Searle AG S. R. (1957) "Matted", a new hair mutant in the house mouse: genetics and morphology. *J Embryol Exp Morphol* **5**.
- Setoguchi R., Hori S., Takahashi T. and Sakaguchi S. (2005) Homeostatic maintenance of natural Foxp3(+) CD25(+) CD4(+) regulatory T cells by interleukin (IL)-2 and induction of autoimmune disease by IL-2 neutralization. *J Exp Med* **201**, 723-35.
- Shy M. E., Arroyo E., Sladky J., Menichella D., Jiang H., Xu W., Kamholz J. and Scherer S. S. (1997) Heterozygous P0 knockout mice develop a peripheral neuropathy that resembles chronic inflammatory demyelinating polyneuropathy (CIDP). *J Neuropathol Exp Neurol* **56**, 811-21.
- Silver S. J. and Rebay I. (2005) Signaling circuitries in development: insights from the retinal determination gene

- network. *Development* **132**, 3-13.
- Smith D. K. and Xue H. (1997) Sequence profiles of immunoglobulin and immunoglobulin-like domains. *J Mol Biol* **274**, 530-45.
- Stevens R., Goble C. A. and Bechhofer S. (2000) Ontology-based knowledge representation for bioinformatics. *Brief Bioinform* **1**, 398-414.
- Suzuki K., Hu D., Bustos T., Zlotogora J., Richieri-Costa A., Helms J. A. and Spritz R. A. (2000) Mutations of PVRL1, encoding a cell-cell adhesion molecule/herpesvirus receptor, in cleft lip/palate-ectodermal dysplasia. *Nat Genet* **25**, 427-30.
- Sweet HO O. S., Taylor BA, Rowe L, Davisson MT, Cook S et al. (1990) Rough fur (ruf). *Mouse Genome* **86**, 236-237.
- Talora C., Campese A. F., Bellavia D., Felli M. P., Vacca A., Gulino A. and Screpanti I. (2008) Notch signaling and diseases: an evolutionary journey from a simple beginning to complex outcomes. *Biochim Biophys Acta* **1782**, 489-97.
- Teichmann S. A. and Chothia C. (2000) Immunoglobulin superfamily proteins in *Caenorhabditis elegans*. *J Mol Biol* **296**, 1367-83.
- TMHMM. <http://www.cbs.dtu.dk/services/TMHMM/>.
- Traver D., Herbomel P., Patton E. E., Murphey R. D., Yoder J. A., Litman G. W., Catic A., Amemiya C. T., Zon L. I. and Trede N. S. (2003) The zebrafish as a model organism to study development of the immune system. *Adv Immunol* **81**, 253-330.
- Tydell C. C., David-Fung E. S., Moore J. E., Rowen L., Taghon T. and Rothenberg E. V. (2007) Molecular dissection of prethymic progenitor entry into the T lymphocyte developmental pathway. *J Immunol* **179**, 421-38.
- Varas A., Hager-Theodorides A. L., Sacedon R., Vicente A., Zapata A. G. and Crompton T. (2003) The role of morphogens in T-cell development. *Trends Immunol* **24**, 197-206.
- Waikel R. L., Kawachi Y., Waikel P. A., Wang X. J. and Roop D. R. (2001) Deregulated expression of c-Myc depletes epidermal stem cells. *Nat Genet* **28**, 165-8.
- Wakae K., Magor B. G., Saunders H., Nagaoka H., Kawamura A., Kinoshita K., Honjo T. and Muramatsu M. (2006) Evolution of class switch recombination function in fish activation-induced cytidine deaminase, AID. *Int Immunol* **18**, 41-7.
- Wang S., Krinks M., Lin K., Luyten F. P. and Moos M., Jr. (1997) Frzb, a secreted protein expressed in the Spemann organizer, binds and inhibits Wnt-8. *Cell* **88**, 757-66.
- Wehrli M. and Tomlinson A. (1998) Independent regulation of anterior/posterior and equatorial/polar polarity in the *Drosophila* eye; evidence for the involvement of Wnt signaling in the equatorial/polar axis. *Development* **125**, 1421-32.
- Willett C. E., Cherry J. J. and Steiner L. A. (1997) Characterization and expression of the recombination activating genes (rag1 and rag2) of zebrafish. *Immunogenetics* **45**, 394-404.
- Willett C. E., Cortes A., Zuasti A. and Zapata A. G. (1999) Early hematopoiesis and developing lymphoid organs in the zebrafish. *Dev Dyn* **214**, 323-36.
- Wilson A. L., Shen Y. C., Babb-Cledenon S. G., Rostedt J., Liu B., Barald K. F., Marrs J. A. and Liu Q. (2007) Cadherin-4 plays a role in the development of zebrafish cranial ganglia and lateral line system. *Dev Dyn* **236**, 893-902.
- Xue Y., Zhou F., Fu C., Xu Y. and Yao X. (2006) SUMOsp: a web server for sumoylation site prediction. *Nucleic Acids Res* **34**, W254-7.
- Yaniv K., Isogai S., Castranova D., Dye L., Hitomi J. and Weinstein B. M. (2006) Live imaging of lymphatic development in the zebrafish. *Nat Med* **12**, 711-6.
- Yoshinari N., Ishida T., Kudo A. and Kawakami A. (2009) Gene expression and functional analysis of zebrafish larval fin fold regeneration. *Dev Biol* **325**, 71-81.
- Yu M., Kissling S., Freyschmidt-Paul P., Hoffmann R., Shapiro J. and McElwee K. J. (2008) Interleukin-6 cytokine family member oncostatin M is a hair-follicle-expressed factor with hair growth inhibitory properties. *Exp Dermatol* **17**, 12-9.