

Chapter 6

Influence of sociodemographic and socioeconomic features on treatment outcome in RCTs versus daily psychiatric practice

Rosalind van der Lem
Purdey M. Stamsnieder
Nic J.A. van der Wee
Tineke van Veen
Frans G. Zitman

ABSTRACT

Purpose: Sociodemographic and socioeconomic characteristics of participants in antidepressant and psychotherapy efficacy trials (AETs and PETs) for major depressive disorder (MDD) may limit the generalizability of the results. We compared trial participants to daily practice patients. We subsequently assessed the influence of sociodemographic and socioeconomic status on treatment outcome in daily practice.

Methods: Data on daily practice patients were derived through Routine Outcome Monitoring (ROM). We included 626 patients with MDD according to the MINIplus. Distributions of age, gender, race, marital status and employment status were compared to participants in 63 selected AETs and PETs. Influence of these features on treatment outcome was explored through multivariate regression analysis.

Results: Trial participants were older, more often male (diff. 4%, $p=0.05$), white (diff 4%, $p<0.001$) and not married (diff 7%, $p=0.003$). Although significant, most differences were relatively small. However, the difference in employment status was striking: 34% of the ROM patients were currently working, versus 68% of the trial participants (diff. 34%, $p<0.001$). Being employed contributed to a positive treatment outcome: OR 1.8 for response (50% reduction of MADRS), OR 1.9 for remission ($MADRS \leq 10$).

Conclusions: employment status should be taken into account while interpreting results from RCTs and as predictor of treatment success in daily practice.

Keywords: major depressive disorder, randomized controlled trial, sociodemographic status, socioeconomic status, patient selection

INTRODUCTION

International guidelines on treatment of major depressive disorder are based on evidence retrieved from scientific research. Meta-analyses and large randomized controlled trials (RCTs) are considered to give the most reliable evidence on treatment outcome for most therapies. RCTs use very strict procedures in design, patient selection, randomization, and methodology to enhance internal validity, i.e. the ability to adequately determine the efficacy of an intervention versus a control condition. While doing so, RCTs diminish their external validity, i.e. the generalizability to routine daily practice. Therefore, many clinicians and researchers have stated that the practical value of RCTs as source of evidence for treatment efficacy in daily practice is limited [1-7].

In previous research [1], many differences which might influence treatment outcome between trial methodology and daily practice have been described. One of the important domains that influence external validity is the selection of participants. The methodology of recruitment, the use of eligibility criteria, the (unintentional) use of criteria beyond eligibility (e.g. by recruitment only of patients with medical insurance, or in a certain area), the use of run-in periods and enrichment strategies, all lead to the exclusion of possible non-responders, and may lead to selection bias in the research population. For major depressive disorder (MDD), most research has focused on the influence of eligibility criteria on external validity. Several researchers [8-11] have shown that only a minority of daily practice patients would be eligible for RCTs. The influence of these eligibility criteria on treatment outcome is not yet clear: Wisniewski and colleagues [9] found that eligible patients had better treatment outcome than non-eligible patients, on the other hand, in our own research [11], we found no differences in treatment outcome between eligible and non-eligible patients. However, the use of eligibility criteria is only one aspect of patient selection. Even if eligibility criteria are met, recruited RCT participants differ from not recruited participants in terms of age, sex, race, educational status, social class and place of residence [1]. In previous research, we found a great variety in the reporting of sociodemographic and socioeconomic (SES) features in major depression trials. SES factors are known to influence treatment outcome in MDD [12-22]. Differences in SES status of participants might be one of the explanations for the differences between treatment outcome in RCTs and in pragmatic trials. However, although much effort is put in approaching daily practice as much as possible in pragmatic trials like STAR*D, they still have several properties of an RCT and selection bias by recruitment is not completely ruled out [23].

In the current project, we first compared the SES status of participants in psychotherapy efficacy trials (PETs) and antidepressants efficacy trials (AETs) to the SES status of a daily practice population of patients suffering from MDD. Subsequently, we explored the influence of the SES status on treatment outcome in daily practice to assess whether possible SES differences between RCT participants and daily practice patients contribute to differences

in treatment outcome between RCTs and routine practice. Comprehensive data on patients' characteristics, treatment modalities and treatment outcome were available through an extensive Routine Outcome Monitoring system (ROM) and extensive chart review. ROM provides outcome data on treatments without the properties of clinical trial but with strong methodological features. Therefore ROM is a qualified instrument to obtain insight in "real life daily practice" [24].

Aims of the study

We investigate the differences in sociodemographic and socioeconomic features between "real life" patients suffering from MDD and trial participants. Furthermore we explore the influence of sociodemographic and socioeconomic features on treatment outcome. This study gives insight on the extent of generalizability of results from RCTs to daily practice as well as the influence of sociodemographic and socioeconomic status on treatment outcome in daily practice.

METHODS

Selection of RCTs and definitions of SES features in RCTs

We included peer reviewed publications of RCTs, published through 2007 in outpatients with a unipolar, non-psychotic major depressive disorder according to DSM-III-R or DSM-IV. Because we aimed to review the reporting of sociodemographic and socioeconomic features in RCTs usually selected for the development of guidelines for routine treatment, we excluded trials which a priori included only participants from specified subgroups like elderly or a specific ethnic minority. Furthermore, it was essential that the publication provided baseline information on sociodemographic and/or socioeconomic features. We included trials written in English, since international guidelines for treatment of MDD are predominantly based on English literature. Because of the large number of published AETs, we restricted our search to AETs published in journals from the top ten Impact Ranking psychiatric journals of 2005. By including only high impact factor journals, we expected to have a sample of trials with the most systematic manner of reporting SES features. For PETs, which are less frequently published, we performed a Medline search for RCTs investigating psychotherapy (cognitive behavior therapy and interpersonal therapy) for adult patients suffering from MDD. We performed an additional search in PsycInfo and checked the reference lists of included trials for other relevant studies as well as the database <http://www.psychotherapyrcts.org>. 45 AETs and 19 PETs were selected. Detailed information on RCT-selection and exploration of SES features of RCT participants has been previously reported [25]. The selected AETs and PETs were equally distributed between Europe and the United States of America. Of all the different SES features mentioned in the selected RCTs,

we only used the SES features that were reported in at least 15% of the selected trials (n=10). In RCTs, SES features are described in various manners. In order to allow an estimation of the mean socioeconomic status of RCT participants, the reporting of the SES feature in the RCT had to be in such manner that the feature could be converted in a dichotomous variable. The following SES features were reported frequently enough (in at least 15% of the trials) and were suitable for comparison between RCTs and our ROM population: age, gender, race, marital status and employment status. Educational level was relatively often reported in RCTs but in so many different ways that dichotomization was not possible. Income and social position were reported in less than 15% of the RCTs. The selected SES features were defined as follows:

- Age: age at baseline assessment calculated in years.
- Gender: percentage females in the population.
- Race: percentage white patients in the population.
- Marital status: percentage of married/cohabitating patients.
- Employment status: percentage of patients who have a paid job.

Treatment outcome in MDD trials is usually defined as percentage of responders (50% reduction of symptoms) or as percentage of remitters (score below a certain severity cut-off).

Patient selection and definitions of SES features in the Routine Outcome Monitoring population

In the Netherlands, health insurance for all citizens is regulated by the government. Therefore, (mental) health care is easily accessible and not restricted by the financial means of individual patients. The Dutch health care system is organised in a stepped-care-manner and described in treatment guidelines. First, patients visit their general practitioner (GP). In case of depression, the treatment guidelines for GPs recommend that patients with mild depression are treated by their GP with lifestyle advices. If the depression is moderate (clinical judgment) the GP can decide to prescribe antidepressants. Reasons to refer patients to a regional mental health provider (RMHP) are: preference of patients for psychotherapy (not provided by GPs); severe depression (clinical judgment); presence of co morbid disorders; complex situation of the patient due to physical health problems; social problems etc.; duration of the depression; and lack of result of antidepressant therapy. After baseline assessment and a clinical interview at our RMHP, patients suffering from major depression are offered to choose between psychotherapy and antidepressant therapy (if the severity is not too high, judged by the interviewing clinician in combination with results from ROM). When patients are already taking antidepressants from their GP, if needed the dose is optimized, and subsequently patients are offered to switch to another antidepressant or psychotherapy is added to the antidepressant therapy.

In 2002, the Regional Mental Health Provider (RMHP) Rivierduinen (service area with 1.1 million inhabitants), in collaboration with the University Medical Hospital Leiden, implemented a comprehensive system of routine outcome monitoring (ROM) and evidence based, stepped care protocols. In ROM, all patients referred to the RMHP for treatment of a mood, anxiety or somatoform disorder have an extensive baseline assessment. Treatment progress is then assessed at intervals of three to four months and before starting a new treatment step. The baseline assessment comprises a standardized diagnostic interview (Mini-International Neuropsychiatric Interview Plus [26]), the collection of sociodemographic and socioeconomic data, the administration of observer rated scales and self report questionnaires, and general measures of health and quality of life. All patients with sufficient mastery of the Dutch language who are able to complete computerized and written questionnaires are eligible for ROM. After baseline assessment and a clinical interview, patients suffering from major depression are offered to choose between psychotherapy and antidepressant therapy. When ROM data are used for research purposes, these are provided to researchers in an anonymous form, as dictated by the Psychiatric Academic Registration Leiden (PAREL) regulation. This procedure has been approved by the Medical Ethical Committee of the University Medical Hospital Leiden. The design of Routine Outcome Monitoring has been reported previously [27].

From the ROM population who sought treatment at the RMHP Rivierduinen from January 2002 until January 2007, we included all outpatients who met the following criteria:

- Patients had a DSM-IV diagnosis of a current major depressive disorder as established by the Mini International Neuropsychiatric Interview.
- Patients had at least one follow-up assessment in ROM.
- Patients were not suspected of bipolarity/psychotic features by their clinician.
- Patients were not admitted to an inpatient clinic during follow-up.
- Patients did not have a MADRS-score ≤ 10 at baseline assessment.
- The time-span between baseline and follow-up assessment was not too short (less than four weeks) or too long (more than one year) to provide reliable information.

We conducted an extensive chart review in order to obtain information on the type of offered treatment. We examined possible selection bias due to our inclusion criteria by comparing the baseline severity of the depression and the SES features of the selected patients to the characteristics of patients who had no follow up assessments in ROM. We compared the selected patients to all patients who suffered from major depression according to the MINIplus (including patients who dropped out of treatment after baseline assessment and patients of whom it was not clear if they dropped out or not). Subsequently, we compared the patients with follow-up to patients who did receive treatment, but had no follow-up assessments in ROM. Information on how many patients did receive treatment was obtained through an anonymized database from the medical administration of RMHP Rivierduinen.

Patients in our population received different types of treatment for their depression. Most of the treatments (84%) were in line with international treatment guidelines: antidepressants (19% with additional social supportive therapy/counseling), individual psychotherapy (mostly cognitive behavioral therapy or interpersonal therapy) or combination therapy of pharmacotherapy and cognitive behavioral therapy or interpersonal therapy. Sixteen percent of the patients received treatment other than antidepressants, individual psychotherapy or combination treatment. In line with RCTs we defined treatment-response as a 50% reduction of symptoms on the Montgomery Asberg Rating Scale for Depression (MADRS, [28]) and remission as MADRS \leq 10.

In ROM extensive information on sociodemographic and socioeconomic status is gathered. Age and gender are recorded. Ethnicity is primarily assessed by the recording of patients' and their parents' countries of birth. Consistent with most RCTs, we considered patients who were born (or whose parents were born) in Suriname, Antilles, Turkey and Morocco as non-white. Two-third of the patients from other non-Dutch origins was non-white. We considered patients who were single, living-apart-together, divorced or widowed as "not married/cohabitating".

Like other member states of the European Union, the Netherlands have a social security system which provides sickness benefit to people who are temporarily not able to work and disability benefit to those who are not able to work anymore. In the USA, the provision of sickness or disability benefit depends upon the insurance policies. Other countries provide no sickness or disability benefit or use their own system. Clearly, there is an important difference in financial status between receiving a sickness or disability benefit or not. It is unknown whether in RCTs, patients who receive sickness benefit are considered as having a paid job or not. Therefore, to allow comparison with RCTs, we defined employment status in ROM with two separate variables: employment status I and II:

For employment status I patients who reported having a paid job at baseline assessment are considered as having "paid work". In this definition "paid work" means that patients are *working* at the time of baseline assessment. Patients on sickness or disability benefit, unemployed, student or housewife were considered as having "no paid work". For employment status II patients who receive sickness or disability benefit were classified as having "paid work" while unemployed patients, students and housewives were still considered as having "no paid work". In this definition having "paid work" means that patients receive an income out of a job, but that a substantial part of these patients (53% of the patients classified as having paid work) is not working but on sickness benefit.

Statistical Analysis

We first compared the included sample from our ROM population with the lost-to-follow-up in ROM patients by using Chi-square tests and independent sample t-tests. Subsequently, the SES features of RCT participants were compared to those of our ROM population by

comparison of proportions of two independent groups [29]. The selected RCTs did not report enough information on the characteristic "Age" to allow a comparison between RCT population and our population.

Finally, we examined the influence of SES features on outcome in routine clinical practice. In our sample, there were missing values for the following variables: type of offered treatment (n=28), and socioeconomic characteristics (n=82). Comparison of complete cases and cases with missing data showed differences on several variables such as age, gender and treatment outcome. In such instances, complete case analysis may yield biased estimates [30]. Therefore, the MICE (multivariate imputation by chained equations [31]) method was used to estimate missing values for type of offered treatment, and socioeconomic characteristics. The influence of the sociodemographic (age, gender, race) and socioeconomic (marital status, employment status) on treatment outcome was computed by logistic regression after MICE. Univariate regression models (unadjusted analysis) were used to explore the influence of the individual SES features on treatment outcome. A multivariate regression model (adjusted analysis) was used to explore the joint influence of all SES features on treatment outcome. In this model, type of offered treatment and educational level were entered as possible confounders. In our previous research [11] on the same patient population, we showed that typical clinical features of MDD patients in daily practice, namely co morbid Axis I disorders, substance abuse, and suicidality, which are often used as exclusion criteria in RCTs, were not associated with treatment outcome in clinical practice. A low baseline severity, however, was associated with a less favorable treatment outcome. Therefore, we entered baseline severity of the depression as another possible confounder in our multivariate regression model.

Odds-ratios (OR) and their confidence-intervals were computed by using the robust standard error. Statistical analyses were performed with SPSS 16.0 and STATA10.0.

RESULTS

Daily practice (ROM) population

4157 outpatients were assessed at baseline between January 2002 and January 2007. Of these patients, 1653 suffered from a current major depressive disorder according to the MINIplus. Since in scientific literature it is not well defined when a depression should be considered the "primary diagnosis" or when a so called "co morbid disorder", we included all patients who suffered from major depression according to the MINIplus. From the 1653 patients suffering from major depression, 46% (n=774) had at least one follow-up assessment. Extensive chart-review was done for those 774 patients. 148 patients had to be excluded from further follow-up analysis due to suspected bipolarity/psychotic features, admission to an inpatient clinic during follow-up, remission on the MADRS at baseline or an insufficient

time-span between baseline and follow-up assessment. Finally, 626 patients were eligible for follow-up analysis. Of 76% of these 626 patients, the clinician stated that depressive disorder was the primary clinical diagnosis. 24% of the patients suffered from depression, but it was considered to be a co morbid disorder. Selection of the patient population is described in figure 1. The characteristics of the patient population are described in table 1.




Figure 1. Selection of the follow-up group.

Table 1. Baseline characteristics of ROM patients.

N=626		
Mean age in years (standard deviation)	39.2 (11.4)	
Gender	Male 3	3.7%
	Female	66.3%
% Female¹	66 %	
Presence of co morbid other Axis I disorder	69.9%	(majority anxiety disorder)
Mean MADRS at baseline	25.8 (6.5)	
Type of offered treatment ²	AD	13.8%
	IP	28.4%
	AD+IP	27.9%
	AD+SST	15.9%
	Other	14.1%

Duration of treatment in weeks (95% CI ³)	AD	20.8 (18.7-22.9)
	IP	20.1 (18.5-21.6)
	AD+IP	21.5 (20.0-23.1)
	AD+SST	21.5 (20.0-23.1)
	Other	19.1 (17.1-21.1)
Marital status	Married/cohabitating	52.2%
	Divorced/widowed	16.2%
	Not married	31.6%
% Married/cohabitating¹	52%	
Birth Country	Netherlands	84.9%
	Turkey/Morocco	5.5%
	Suriname/Antilles	2.9%
	Other	6.6%
Parental Birth Country	Netherlands	79.2%
	Turkey/Morocco	7.2%
	Suriname/Antilles	2.9%
	Other	10.7%
Race (% whites)¹	84%	
Employment status	Employed	26.1%
	Not employed	34.4%
	Sickness benefit	39.5%
% paid work I¹	34%	
% paid work II¹	74%	

¹ Dichotomization of variable for comparison to RCT/AET/PET, see Method section.

² AD=antidepressants, IP=individual psychotherapy, SST=social supportive therapy.

³ 95% CI= 95% confidence interval.

The selected patient group was similar to the not-selected patients in baseline severity of the depression and most SES features. The selected patients were slightly older (39.2 versus 37.3 years) and more often married/cohabitating (52.6% versus 46.9%) than the patients who were not selected (see table 2, supplementary material). Of the patients who were lost to follow-up, and therefore not selected, 63% received treatment in our outpatient clinics for mood-, anxiety, and somatoform disorders after baseline assessment (37% was referred to other specialized departments or did drop out of treatment). We also compared the selected patients to the patients who did receive treatment in our outpatient clinics but had no follow-up in ROM and found them to be very similar (see table 3, supplementary material). Therefore we consider our selected patient sample as fairly representative for patients who receive outpatient treatment for depressive disorders.

RCT participants versus daily practice (ROM) population

We compared the RCT population and our ROM sample on age, gender, ethnicity, marital status, and employment status. RCT participants were less often female (62% vs. 66%), more often white (89% vs. 84%), less often married/cohabitating (45% vs. 52%) and more often employed (68% vs. 34%) than daily practice (ROM) patients. Both in antidepressant and psychotherapy trials, participants were more often white, less often married/cohabitating and more often employed than daily practice (ROM) patients. Both in antidepressant and psychotherapy trials, the male-female ratio was different than in daily practice (ROM): in antidepressant trials the ratio is in favor of males (39% vs. 34% male) compared to daily practice (ROM), while in psychotherapy trials the ratio is in favor of females (73% vs. 66% female) compared to daily practice (ROM). Although the differences were statistically significant, they were sometimes relatively small and significance was probably reached due to the large number of patients included in both groups. However, the exception is the difference between RCT participants and our daily practice population in employment status (employment status I: defined as currently having “paid work”, not on sickness or disability benefit); 68% of the RCT participants were currently employed, whereas only 34% of the daily practice patients had a paid job. Results from the comparison between RCT participants and our population are described in table 4.

Table 4. SES features of RCT participants versus ROM patients.

	RCT (%, n)	AET (%, n)	PET (%, n)	ROM ² (%, n)	Difference in proportion RCT versus ROM (95% CI; p-value)	Difference in proportion AET versus ROM (95% CI, p-value)	Difference in proportion PET versus ROM (95% CI, p-value)
Age in Years¹	41	41	37	39 (sd 11.4)	-	-	-
Gender (% female)	62% (5880)	61% (5269)	73% (611)	66% (415)	-4% (-8% – 0%; 0.05)	-5% (-9% – -11%; 0.016)	7% (2% – 11%; 0.008)
Race (% white)	89% (5820)	89% (5400)	85% (420)	84% (457)	4% (2% – 8%; <0.001)	4% (2 – 8%; <0.001)	-1% (-3% – -6%; NS)
Marital status (% married)	45% (972)	46% (728)	42% (244)	52% (284)	-7% (-12% – -3%; 0.003)	-10% (-10% – -1%; 0.017)	-18% (-24% – -12%; <0.001)
Employment I³ (% paid work)	68% (1107)	68% (905)	66% (202)	34% (187)	34% (29% – 38%; p<0.001)	34% (30% – 39%; <0.001)	32% (25% – 38%; <0.001)
Employment II (% paid work or sickness benefit)	68% (1107)	68% (905)	66% (202)	74% (402)	-6% (-10% – -1%; 0.013)	-5% (-10% – -1%; 0.027)	-8% (-14% – -1%; 0.019)

¹ Statistical comparison not possible.

² For all 626 patients in ROM information on age and gender was available, for 82 patients no information was available on race, marital status and employment status.

³ In Employment I, paid work is defined as only patients who are currently working, Employment II includes both patients who are currently working and who receive a sickness benefit.

AET: Antidepressant Efficacy Trial; PET: Psychotherapy Efficacy Trial; ROM: data from routine outcome monitoring; CI: confidence Interval; n: total number of patients studied in the included RCTs/AETs/PETs or ROM.

Influence of SES features on treatment outcome in the daily practice (ROM) population

In the univariate analyses, having a paid job (employment status I, in which patients who receive sickness benefit were defined as not having a paid job) contributed positively to remission ($MADRS \leq 10$ [28]) and response (50% reduction of the MADRS). If patients on sickness benefit are also considered to have a paid job (employment status II), the influence of employment status on treatment outcome disappeared. None of the other SES features contributed significantly to treatment outcome. In the multivariate analyses, we investigated the influence of Age, Gender, Race, Marital status, and Employment status (Employment I) on treatment outcome in one model. In this model we also adjusted for baseline severity of the depression, so that we could analyze the influence of SES features irrespective of the severity of the illness. We found that having a paid job contributed positively to remission (OR 1.85, 95%CI 1.2–2.8; RSE 0.21; $p=0.003$, R-square 7%), and response (OR 1.76, 95%CI 1.2–2.6; RSE 0.20; $p=0.005$, R-square 3%). Results from the unadjusted analysis of the influence of the individual SES features are described in table 5; Results from the analysis of the joint influence of all SES features adjusted for baseline severity of the depression, type of offered treatment and educational level are described in table 6.

Table 5. Influence of individual SES features on treatment outcome: unadjusted analysis.

	Remission (MADRS ≤ 10)	Response (50% reduction MADRS)
Age	OR 0.99 (95%CI 0.97–1.00; RSE 0.008; $p=0.13$)	OR 0.99 (95%CI 1.0v1.0; RSE 0.01; $p=0.10$)
Gender	OR 0.97 (95%CI 0.65–1.44; RSE 0.20; $p=0.87$)	OR 1.09 (95%CI 0.8–1.6; RSE 0.21; $p=0.66$)
Race	OR 0.48 (95%CI 0.27–1.1; RSE 0.41; $p=0.08$)	OR 0.51 (95%CI 0.2–1.1; RSE 0.41; $p=0.08$)
Marital status	OR 1.32 (95%CI 0.9–2.0; RSE 0.02; $p=0.17$)	OR 1.32 (95% CI 0.9–2.0; RSE 0.20; $p=0.17$)
Employment I	OR 2.30 (95%CI 1.6–3.4; RSE 0.20; $p<0.001$)	OR 1.82 (95%CI 1.3–2.6; RSE 0.34; $p=0.002$)
Employment II	OR 1.12 (95%CI 0.7–1.1; RSE 0.23; $p=0.61$)	OR 1.18 (95%CI 0.8–1.8; RSE 0.21; $p=0.42$)

OR: odds ratio

CI: confidence interval

RSE: robust standard error

Table 6. Joint influence of SES features on treatment outcome: adjusted analysis. Age, gender, race, marital status and employment I status entered in one model as predictors of outcome corrected for severity of depression at baseline, type of offered treatment and educational level.

	Remission (MADRS \leq10)	Response (50% reduction of MADRS)
Age	OR 0.99 (95%CI 0.97–1.00; RSE 0.01; p=0.21)	OR 0.99 (95%CI 0.97–1.00; RSE 0.01; p=0.15)
Gender	OR 1.07 (95%CI 0.69–1.67; RSE 0.23; p=0.75)	OR 1.10 (95%CI 0.73–1.59; RSE 0.20; p=0.65)
Race	OR 0.67 (95%CI 0.32–1.40; RSE 0.38; p=0.29)	OR 0.63 (95%CI 0.33–1.19; RSE 0.32; p=0.15)
Marital status	OR 1.33 (95%CI 0.85–2.07; RSE 0.23; p=0.21)	OR 1.10 (95%CI 0.73–1.67; RSE 0.21; p=0.64)
Employment I	OR 1.89 (95%CI 1.24–2.81; RSE 0.21; p=0.003)	OR 1.76 (95%CI 1.19–2.60; RSE 0.20; p=0.004)

OR: odds ratio

CI: confidence interval

RSE: robust standard error

DISCUSSION

In the current project, we compared sociodemographic and socioeconomic features of participants in randomized controlled trials for depression to those of patients in daily practice. Participants in RCTs for major depression differed from daily practice patients with respect to age, male-female ratio, ethnicity and marital status, but those differences were relatively small (less than 7% difference on all features). One striking difference between RCT participants and daily practice patients is their employment status. Only 34% of the daily practice patients had a paid job (patients who were students, housewives, unemployed or on sickness or disability benefit were considered as having “no paid work”) at time of assessment, while 68% of the RCT participants had a paid job at time of the trial participation. In routine clinical practice, having a paid job contributed positively to treatment outcome, both on remission (OR 1.85) and response (OR 1.76) on the MADRS. Age, gender, race and marital status did not contribute to treatment outcome in routine clinical practice.

Previous research found unequivocal results on the associations between sociodemographic and socioeconomic features and treatment outcome: in general, increased age seemed to be associated with worse outcome in pharmacotherapy [13-19], but not in psychotherapy [20-22], and one study found that men had better treatment outcome in psychotherapy than women [21]. Patients who were married benefitted more from both pharmacotherapy and psychotherapy [18-21,32-35].

The difference in employment status between RCT participants and daily-practice patients was striking. In a recent study was found that the quality of mental health is associated with lesser years of unemployment [36]. In previous research, in non-routine practice settings, on the relation between employment and treatment success, it was found that employment status improved outcome for antidepressants, but not for psychotherapy [18-21,32-35]. In the univariate analyses we found that “currently working” (employment status I) is associated with better treatment outcome, while “receiving income out of a job” (employment status II) seemed not be associated with treatment success. It is, however, likely that other clinical factors, such as baseline severity, could have acted as a confounder in these univariate analyses, since patients with a more severe depression might drop out from work earlier. Therefore, in our multivariate analyses we adjusted for baseline severity of the depression. Still, having paid work (defined as currently working) almost doubled the probability of response or remission. From clinical experience, one might expect that the social status and the daily structure, routine and distraction provided by paid work will be factors that contribute to the positive response on depression treatment. Alternatively, it may be that patients who remain working despite the opportunity of sickness benefit have personality traits that increase the likelihood of positive treatment outcome, like for example optimism. Future research is needed to explore these different possibilities.

Of course, besides differences in SES features, there are many other differences between RCT participants and daily practice patients, caused by the use of stringent exclusion criteria in RCTs. For instance, the presence of co morbid Axis I disorders is used as an exclusion criterion in more than 75% of the antidepressant efficacy trials [37] and in 25–50% of the psychotherapy efficacy trials [38]. In previous research, we found that only 17–24% of our patients would be eligible for participation in antidepressant efficacy trials [11]. Most of our patients would have been excluded because of the presence of co morbid Axis I disorders and not meeting a baseline severity threshold. However, in our previous research [11], we also found that in daily practice the exclusion of patients with co morbid Axis I disorders does not influence the treatment result for depression. One can therefore argue that it is unlikely that co morbidity acted as a major confounder in our analysis of the influence of SES features on treatment outcome for depression. It is, however, very well possible that having a co morbid disorder could lead to earlier withdrawing from work. Since co morbid disorders occur frequently in depressed patients, more specific research on the association between drop-out from work and co morbidity is recommended.

Strengths

We consider the large sample of well-characterized, routinely monitored patients from routine clinical practice to be the major strength of our study. To our knowledge, no previous research has reported on the influence of sociodemographic and socioeconomic features on treatment outcome in major depression in routine daily clinical practice.

Limitations

There are also some limitations to consider. In our model of sociodemographic and socioeconomic features of RCT participants, we converted all information on these features reported in the individual RCTs into dichotomous or trichotomous variables. This conversion led to loss of information, but was necessary for comparability with published data [25]. Another limiting factor might be that in the model AETs outnumbered PETs, both in number of included trials and number of included patients per trial. Thus, the features of the RCT-participants as a group were dominated by the features of AET participants [25]. There was a considerable loss-to-follow up in our sample. Nevertheless, we demonstrated that our patient selection was fairly representative for the daily practice patients who received treatment. It was not possible to analyze in our study which aspect of “having a paid job” contributed positively to a favorable treatment outcome. Furthermore, no information on the duration and the number of episodes of depression was available in our ROM data. Chronic depression is known to have a less favorable prognosis and it is possible that especially patients who suffer from chronic depression have to resign from work, which may have confounded our results. Finally, since sociodemographic and socioeconomic features are closely related to the culture in the country of origin in which research takes place, cultural aspects might have somewhat limited the generalizability of our results. It is unknown to what extent the Dutch health care system (and that of several other, mostly European, countries) may limit the generalizability of our results to other countries that do not have an extensive social security system. Furthermore, we have to take into account that we compared a western psychiatric population to an RCT population derived from western countries (Europe and USA). Since our aim was to compare RCT participants (who are most often from western countries) to our daily practice patients, the fact that all RCT participants were from western countries did not limit our research. However, our results are probably not generalizable to countries outside the western world. Future research on the generalizability of results from RCTs to psychiatric patients in other parts of the world is highly recommended.

CONCLUSION

In conclusion, we found that RCT participants and daily practice patients only differed slightly on most sociodemographic and socioeconomic features, with the exception of having a paid job. Having a paid job contributed significantly to treatment success in daily practice and should be taken into account both while interpreting results from RCTs as well as in depression treatment in daily practice. Further research is recommended to explore which specific aspects of employment status contribute to better treatment outcome.

REFERENCE LIST

1. Rothwell PM: External validity of randomised controlled trials: "to whom do the results of this trial apply?". *Lancet* 2005, 365: 82-93.
2. Wells KB: Treatment research at the crossroads: the scientific interface of clinical trials and effectiveness research. *Am J Psychiatry* 1999, 156: 5-10.
3. Tunis SR, Stryer DB, Clancy CM: Practical clinical trials: increasing the value of clinical research for decision making in clinical and health policy. *JAMA* 2003, 290: 1624-1632.
4. Mulder RT, Frampton C, Joyce PR, Porter R: Randomized controlled trials in psychiatry. Part II: their relationship to clinical practice. *Aust N Z J Psychiatry* 2003, 37: 265-269.
5. Licht RW, Gouliaev G, Vestergaard P, Frydenberg M: Generalisability of results from randomised drug trials. A trial on antimanic treatment. *Br J Psychiatry* 1997, 170:264-7.
6. Stewart JW, McGrath PJ, Quitkin FM: Can mildly depressed outpatients with atypical depression benefit from antidepressants? *Am J Psychiatry* 1992, 149: 615-619.
7. Moller HJ: How close is evidence to truth in evidence-based treatment of mental disorders? *Eur Arch Psychiatry Clin Neurosci* 2011.
8. Zimmerman M, Mattia JI, Posternak MA: Are subjects in pharmacological treatment trials of depression representative of patients in routine clinical practice? *Am J Psychiatry* 2002, 159: 469-473.
9. Wisniewski SR, Rush AJ, Nierenberg AA, Gaynes BN, Warden D, Luther JF *et al.*: Can phase III trial results of antidepressant medications be generalized to clinical practice? A STAR*D report. *Am J Psychiatry* 2009, 166: 599-607.
10. Zetin M, Hoepner CT: Relevance of exclusion criteria in antidepressant clinical trials: a replication study. *J Clin Psychopharmacol* 2007, 27: 295-301.
11. van der Lem R, van der Wee NJ, van Veen T., Zitman FG: The generalizability of antidepressant efficacy trials to routine psychiatric out-patient practice. *Psychol Med* 2011, 41: 1353-1363.
12. Button K.S., Wiles N.J., Lewis G, Peters T.J., Kessler D.: Factors associated with differential response to online cognitive behavioural therapy. *Soc Psychiatry Psychiatr Epidemiol* 2012, 47: 827-833.
13. Aberg-Wistedt A, Agren H, Ekselius L, Bengtsson F, Akerblad AC: Sertraline versus paroxetine in major depression: clinical outcome after six months of continuous therapy. *J Clin Psychopharmacol* 2000, 20: 645-652.
14. Kornstein SG, Schatzberg AF, Thase ME, Yonkers KA, McCullough JP, Keitner GI *et al.*: Gender differences in treatment response to sertraline versus imipramine in chronic depression. *Am J Psychiatry* 2000, 157: 1445-1452.
15. Joyce PR, Mulder RT, Luty SE, Sullivan PF, McKenzie JM, Abbott RM *et al.*: Patterns and predictors of remission, response and recovery in major depression treated with fluoxetine or nortriptyline. *Aust N Z J Psychiatry* 2002, 36: 384-391.
16. Papakostas GI, Petersen T, Mischoulon D, Hughes ME, Spector AR, Alpert JE *et al.*: Functioning and interpersonal relationships as predictors of response in treatment-resistant depression. *Compr Psychiatry* 2003, 44: 44-50.
17. Baca E, Garcia-Garcia M, Porras-Chavarino A: Gender differences in treatment response to sertraline versus imipramine in patients with nonmelancholic depressive disorders. *Prog Neuropsychopharmacol Biol Psychiatry* 2004, 28: 57-65.
18. Lowe B, Schenkel I, Bair MJ, Gobel C: Efficacy, predictors of therapy response, and safety of sertraline in routine clinical practice: prospective, open-label, non-interventional postmarketing surveillance study in 1878 patients. *J Affect Disord* 2005, 87: 271-279.
19. Trivedi MH, Rush AJ, Wisniewski SR, Nierenberg AA, Warden D, Ritz L *et al.*: Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry* 2006, 163: 28-40.
20. Jarrett RB, Eaves GG, Grannemann BD, Rush AJ: Clinical, cognitive, and demographic predictors of response to cognitive therapy for depression: a preliminary report. *Psychiatry Res* 1991, 37: 245-260.

21. Sotsky SM, Glass DR, Shea MT, Pilkonis PA, Collins JF, Elkin I *et al.*: Patient predictors of response to psychotherapy and pharmacotherapy: findings in the NIMH Treatment of Depression Collaborative Research Program. *Am J Psychiatry* 1991, 148: 997-1008.
22. Thase ME, Reynolds CF, III, Frank E, Simons AD, McGeary J, Fasiczka AL *et al.*: Do depressed men and women respond similarly to cognitive behavior therapy? *Am J Psychiatry* 1994, 151: 500-505.
23. Hatcher S: The STAR*D trial: the 300 lb gorilla is in the room, but does it block all the light? *Evid Based Ment Health* 2008, 11: 97-99.
24. Percevic R, Lambert MJ, Kordy H: Computer-supported monitoring of patient treatment response. *J Clin Psychol* 2004, 60: 285-299.
25. Lem Rvd, Stamsnieder P, Wee Nvd, Veen Tv, Zitman FG (Eds): Socio-demographic features in randomized controlled trials for major depression: generalizability and individualization. In *Int J Person Cent Medicine* 2011, 1: 268-278.
26. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E *et al.*: The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 1998, 59 Suppl 20: 22-33.
27. de Beurs E., den Hollander-Gijsman ME, van Rood YR, van der Wee NJ, Giltay EJ, van Noorden MS *et al.*: Routine outcome monitoring in the Netherlands: practical experiences with a web-based strategy for the assessment of treatment outcome in clinical practice. *Clin Psychol Psychother* 2011, 18: 1-12.
28. Asberg M, Montgomery SA, Perris C, Schalling D, Sedvall G: A comprehensive psychopathological rating scale. *Acta Psychiatr Scand Suppl* 1978, 5-27.
29. Altman DG: Practical Statistics for Medical Research. 1991:229-276.
30. Donders AR, van der Heijden GJ, Stijnen T, Moons KG: Review: a gentle introduction to imputation of missing values. *J Clin Epidemiol* 2006, 59: 1087-1091.
31. Royston P: Multiple imputation of missing values: update. *Stata Journal* 2005, 5: 188-201.
32. Hollon SD, Derubeis RJ, Evans MD, Wiemer MJ, Garvey MJ, Grove WM *et al.*: Cognitive therapy and pharmacotherapy for depression. Singly and in combination. *Arch Gen Psychiatry* 1992, 49: 774-781.
33. Falconnier L: Socioeconomic status in the treatment of depression. *Am J Orthopsychiatry* 2009, 79: 148-158.
34. Goekoop JG, Hoeksema T, Knoppert-Van der Klein EA, Klinkhamer RA, Van Gaalen HA, Van Londen L *et al.*: Multidimensional ordering of psychopathology. A factor-analytic study using the Comprehensive Psychopathological Rating Scale. *Acta Psychiatr Scand* 1992, 86: 306-312.
35. Van HL, Schoevers RA, Dekker J: Predicting the outcome of antidepressants and psychotherapy for depression: a qualitative, systematic review. *Harv Rev Psychiatry* 2008, 16: 225-234.
36. Butterworth P, Leach L.S., Pirkis J., Kelaher M: Poor mental health influences risk and duration of unemployment: a prospective study. *Soc Psychiatry Psychiatr Epidemiol* 2012, 47: 1013-1021.
37. Posternak MA, Zimmerman M, Keitner GI, Miller IW: A reevaluation of the exclusion criteria used in antidepressant efficacy trials. *Am J Psychiatry* 2002, 159: 191-200.
38. van der Lem R, de Wever WW, van der Wee NJ, van VT, Cuijpers P, Zitman FG: The generalizability of psychotherapy efficacy trials in major depressive disorder: an analysis of the influence of patient selection in efficacy trials on symptom outcome in daily practice. *BMC Psychiatry* 2012, 12: 192.

SUPPLEMENTARY MATERIAL

Table 2. Comparison of all patients who had baseline assessment (follow-up: treatment + follow-up assessment; lost to follow-up: drop out or treatment + no follow up assessment).

	Baseline severity of depression (MADRS)	Age (years)	Gender (% female)	Race (% white)	Marital status (% married)	Employment I (% paid work)	Employment II (% paid work or sickness benefit)
Selected Patientgroup	25.8	39.2	66.3%	82.0%	52.6%	34.1%	74.4%
Lost to follow-up	25.6	37.3	66.6%	80.5%	46.9%	38.8%	70.8%
Statistical Comparison	CI of difference: -0.90-0.63; p=0.73	CI of difference: 0.71-3.09; p=0.002	X ² =0.02 Df1 p=0.89	X ² =0.65 Df2 p=0.72	X ² =3.47 Df1 p=0.05	X ² =2.82 Df1 p=0.09	X ² =2.01 Df1 p=0.16

For 82 of the selected patients no information was available on race, marital status, and employment status. For 291 patients in the lost-to follow up group no information was available on race, marital status, and employment status.

Table 3. Comparison of patients who had treatment+ follow-up assessment and patient who had treatment without follow-up assessment.

	Baseline severity of depression (MADRS)	Age (years)	Gender (% female)	Race (% white)	Marital status (% married)	Employment I (% paid work)	Employment II (% paid work or sickness benefit)
Selected Patientgroup	25.8	39.2	66.3%	82.0%	52.6%	34.1%	74.4%
Lost to follow-up	25.4	37.7	68.1%	80.6%	47.2%	38.5%	71.8%
Statistical Comparison	CI of difference: -1.19-0.43 p=0.35	CI of difference: 0.30-2.83 p=0.015	X ² =0.49 Df1 p=0.48	X ² =0.29 Df2 p=0.86	X ² =3.09 Df1 p=0.08	X ² =2.15 Df1 p=0.14	X ² =0.96 Df1 p=0.33

For 82 of the selected patients no information was available on race, marital status, and employment status. For 153 patients in the lost-to follow up group no information was available on race, marital status, and employment status.

CI: confidence interval. Df: degrees of freedom. X²: chi square



