

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/32843> holds various files of this Leiden University dissertation

Author: Smit, Renske

Title: Star-forming galaxies at the cosmic dawn = Stervormende sterrenstelsels tijdens het kosmische ochtendgloren

Issue Date: 2015-04-28

Star-Forming Galaxies at the Cosmic Dawn

Renske Smit

Cover: River Lek Sunrise, *credit: Sara Winter*, and MACS J0416.1-2403, *credit: NASA, ESA, and M. Postman (STScI), and the CLASH team.*

ISBN: 978-94-6108-9632

Star-Forming Galaxies at the Cosmic Dawn

Stervormende sterrenstelsels tijdens het kosmische ochtendgloren

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. C.J.J.M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op dinsdag 28 April 2015
klokke 16:15 uur

door

Renske Smit
geboren te Los Angeles
in 1985

Promotiecommissie

Promotor: Prof. dr. Marijn Franx

Co-Promotor: Dr. Rychard Bouwens

Overige leden: Prof. dr. Daniel Schaerer (Université de Geneve)

Prof. dr. Huub Röttgering

Prof. dr. Joop Schaye

dr. Jarle Brinchman

dr. Ivo Labbé

*Hoe kón ik dat niet eerder weten,
niet beter zien in vroeger tijd?
Hoe moet ik het weer ooit vergeten?
– M. Vasalis¹*

¹**Tijd** uit Parken en Woestijnen

Contents

1	Introduction	11
1.1	Early galaxy formation	12
1.2	Observational challenges	13
1.3	Recent results on high-redshift galaxies	15
1.3.1	UV luminosity functions and the cosmic star formation rate density	16
1.3.2	Stellar masses and specific star formation rates	18
1.4	Thesis summary	20
2	The Star Formation Rate Function for Redshift $z \sim 4 - 7$ Galaxies	23
2.1	Introduction	24
2.2	SFR functions	24
2.2.1	Dust-corrected luminosity functions	25
2.2.2	Analytical SFR functions	27
2.3	Results	29
2.3.1	High-SFR end: linear build-up of $\log \text{SFR}^*$ from $z \sim 8$ to $z \sim 2$	29
2.3.2	Low-SFR end: flatter slopes and large uncertainties	31
2.4	Summary	31
3	High-EW Nebular Emission in $z \sim 7$ Galaxies	39
3.1	Introduction	40
3.2	Observations	41
3.2.1	Data	41
3.2.2	Photometry and Selection	42
3.2.3	IRAC Photometry	43
3.3	Results	44
3.3.1	[3.6]–[4.5] color distribution and nebular emission lines	44
3.3.2	Inferred [OIII]+H β EWs of $z \sim 7$ galaxies from the mean SED	46
3.3.3	Specific star formation rates	48
3.4	Summary and discussion	51
4	High-precision Photometric Redshifts from Spitzer/IRAC	55
4.1	Introduction	56
4.2	Observations, photometry and $z \sim 5 - 8$ sample	57
4.2.1	<i>HST</i> data and photometry	57
4.2.2	<i>Spitzer</i> /IRAC data and photometry	58
4.2.3	Base sample of $z \sim 5 - 8$ galaxies	59
4.3	[3.6]–[4.5] color vs. redshift	60
4.4	Ultra-blue [3.6]–[4.5] galaxies	61
4.4.1	Blue [3.6]–[4.5] sources at $z \sim 6.8$	63
4.4.2	Blue [3.6]–[4.5] sources at $z \sim 6.0$	65
4.5	A fiducial sample of $z \sim 6.8$ emission line galaxies	67
4.5.1	Selection of our fiducial $z \sim 6.8$ sample	67

4.5.2	Ascertaining the mean redshift and contamination fraction of the $z \sim 6.8$ sample	68
4.5.3	Quantifying the rest-frame EWs of [OIII]+H β in our $z \sim 6.8$ IRAC ultra-blue sample	70
4.6	Summary and discussion	71
5	On the Systematic Use of Hα to Measure Star Formation Rates	77
5.1	Introduction	78
5.2	Data and Samples	79
5.2.1	Spectroscopic sample	79
5.2.2	Photometric sample	81
5.3	Derived properties of $z = 3.8 - 5.0$ galaxies	82
5.3.1	SED fitting	82
5.3.2	H α + [NII] equivalent widths and line strengths	82
5.3.3	Composite SEDs	83
5.4	Inferred H α as star formation rate indicator	84
5.4.1	Star formation rate indicators	84
5.4.2	Origin of high H α star formation rates	85
5.5	SFR-mass sequence	89
5.6	Star formation rate functions	90
5.7	Summary	91
	Nederlandse samenvatting	97
	Curriculum Vitae	103
	Nawoord	105

