

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/35085> holds various files of this Leiden University dissertation

Author: Fumagalli, Mattia

Title: Star formation and aging at cosmic noon : the spectral evolution of galaxies from $z=2$

Issue Date: 2015-09-08

STAR FORMATION AND AGING
AT COSMIC NOON:
the spectral evolution of galaxies from $z=2$

STAR FORMATION AND AGING
AT COSMIC NOON:
the spectral evolution of galaxies
from $z = 2$

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof.mr. C.J.J.M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op dinsdag 8 september 2015
klokke 10:00 uur

door

Mattia Fumagalli

geboren te Lecco, Italië
in 1986

Promotiecommissie

Promotores: Prof. dr. M. Franx
Prof. dr. P. G. van Dokkum Yale University

Overige leden: Prof. dr. H. J. A. Röttgering
Prof. dr. K. H. Kuijken
Prof. dr. J. Schaye
Prof. dr. M. Kriek University of California at Berkeley
dr. K. I. Caputi Rijksuniversiteit Groningen
dr. I. Labbé

Cover: 3D-HST stacks of spectra of star-forming galaxies pictured as mountain ranges. Peaks fade to the horizon as redshift increases.

Designed by Mattia Fumagalli and Marco Vedoá

To Anna

*Two roads diverged in a wood, and I
I took the one less traveled by
(Robert Frost)*

CONTENTS

1	Introduction	1
1.1	The Birth of Extragalactic Astronomy	1
1.2	The Striking Diversity of Galaxies	1
1.3	Measuring Star Formation through cosmic time	3
1.4	Issues at high redshifts	6
1.5	This Thesis	7
2	Hα Equivalent Widths from the 3D-HST survey	13
2.1	Introduction	14
2.2	Data	15
2.2.1	3D-HST	15
2.2.2	SDSS	15
2.2.3	VVDS	16
2.2.4	High redshift data	16
2.3	The EW(H α) - mass relation	16
2.4	The Evolution of EW(H α) with redshift	18
2.5	The sSFR(H α) - mass relation and its evolution with redshift	18
2.6	Linking the characteristic SFH of galaxies and EW(H α)	21
2.7	Conclusions	22
3	How dead are dead galaxies?	25
3.1	Introduction	26
3.2	Data	26
3.3	Sample selection and motivations of the study	29
3.3.1	Selection of Quiescent Galaxies	29
3.3.2	Spectra and SEDs of the sample	30
3.3.3	SFRs from SED fitting and expectations from gas recycling	31
3.3.4	How much star formation could be hidden?	31
3.4	Measuring Obscured Star-Formation Rates of Quiescent Galaxies	33
3.5	Other possible contributions to L_{IR}	36
3.5.1	AGN	36
3.5.2	Circumstellar dust	36
3.5.3	Cirrus dust	38
3.6	Discussion	38
3.7	Conclusions	40
3.A	Appendix A: Photometry	43
3.B	Appendix B: Field-to-field variation	43

4	Stacked spectra	49
4.1	Introduction	50
4.2	Data	51
4.2.1	The 3D-HST survey	51
4.2.2	Sample Selection	53
4.3	Methods	55
4.3.1	Stacking	55
4.3.2	Model fitting	56
4.4	Quiescent Galaxies	57
4.4.1	Quality of fits	57
4.4.2	Determination of Ages	59
4.5	Star Forming Galaxies	60
4.5.1	Quality of fits	60
4.5.2	Determination of Ages	62
4.6	Discussion	63
4.6.1	Differences among SPSs	63
4.6.2	Evolution of Ages	63
4.6.3	H α in quiescent galaxies	67
4.7	Conclusions	68
5	Decreasing Hα for redder star-forming galaxies: influence of dust and star formation rates	71
5.1	Introduction	72
5.2	Data and Sample Selection	73
5.2.1	The 3D-HST survey	73
5.2.2	Sample selection and the UVJ diagram	74
5.3	EW(H α): trend with color	75
5.3.1	Separating star-forming and quiescent galaxies with the UVJ selection	75
5.3.2	Color dependence of EW(H α) for star-forming galaxies	77
5.4	Specific Star Formation Rates of star-forming galaxies: trend with color	77
5.5	Dust absorption of star-forming galaxies along the UVJ diagram	80
5.5.1	Absorption in H α	80
5.5.2	Absorption of the continuum	83
5.6	Conclusions	87
	Samenvatting	91
	Publications	97
	Curriculum Vitae	103
	Acknowledgments	105

