

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/42085> holds various files of this Leiden University dissertation.

**Author:** Milovic, D.

**Title:** On the 16-rank of class groups of quadratic number fields

**Issue Date:** 2016-07-04

# Bibliography

- [1] Pierre Barrucand and Harvey Cohn. Note on primes of type  $x^2 + 32y^2$ , class number, and residuacity. *J. Reine Angew. Math.*, 238:67–70, 1969.
- [2] Manjul Bhargava. Higher composition laws. I. A new view on Gauss composition, and quadratic generalizations. *Ann. of Math. (2)*, 159(1):217–250, 2004.
- [3] Manjul Bhargava, Arul Shankar, and Jacob Tsimerman. On the Davenport-Heilbronn theorems and second order terms. *Invent. Math.*, 193(2):439–499, 2013.
- [4] H. Cohen and H. W. Lenstra, Jr. Heuristics on class groups of number fields. In *Number theory, Noordwijkerhout 1983 (Noordwijkerhout, 1983)*, volume 1068 of *Lecture Notes in Math.*, pages 33–62. Springer, Berlin, 1984.
- [5] H. Cohn and J. C. Lagarias. On the existence of fields governing the 2-invariants of the classgroup of  $\mathbf{Q}(\sqrt{dp})$  as  $p$  varies. *Math. Comp.*, 41(164):711–730, 1983.
- [6] H. Cohn and J. C. Lagarias. Is there a density for the set of primes  $p$  such that the class number of  $\mathbf{Q}(\sqrt{-p})$  is divisible by 16? In *Topics in classical number theory, Vol. I, II (Budapest, 1981)*, volume 34 of *Colloq. Math. Soc. János Bolyai*, pages 257–280. North-Holland, Amsterdam, 1984.
- [7] Harvey Cohn and George Cooke. Parametric form of an eight class field. *Acta Arith.*, 30(4):367–377, 1976.
- [8] David A. Cox. *Primes of the form  $x^2 + ny^2$* . A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1989. Fermat, class field theory and complex multiplication.
- [9] H. Davenport. On a principle of Lipschitz. *J. London Math. Soc.*, 26:179–183, 1951.
- [10] H. Davenport. Corrigendum: “On a principle of Lipschitz”. *J. London Math. Soc.*, 39:580, 1964.
- [11] H. Davenport and H. Heilbronn. On the density of discriminants of cubic fields. II. *Proc. Roy. Soc. London Ser. A*, 322(1551):405–420, 1971.
- [12] Etienne Fouvry and Henryk Iwaniec. Gaussian primes. *Acta Arith.*, 79(3):249–287, 1997.
- [13] Étienne Fouvry and Jürgen Klüners. Cohen-Lenstra heuristics of quadratic number fields. In *Algorithmic number theory*, volume 4076 of *Lecture Notes in Comput. Sci.*, pages 40–55. Springer, Berlin, 2006.

- [14] Étienne Fouvry and Jürgen Klüners. On the 4-rank of class groups of quadratic number fields. *Invent. Math.*, 167(3):455–513, 2007.
- [15] Étienne Fouvry and Jürgen Klüners. On the negative Pell equation. *Ann. of Math. (2)*, 172(3):2035–2104, 2010.
- [16] Étienne Fouvry and Jürgen Klüners. The parity of the period of the continued fraction of  $\sqrt{d}$ . *Proc. Lond. Math. Soc. (3)*, 101(2):337–391, 2010.
- [17] J. B. Friedlander, H. Iwaniec, B. Mazur, and K. Rubin. The spin of prime ideals. *Invent. Math.*, 193(3):697–749, 2013.
- [18] John Friedlander and Henryk Iwaniec. Asymptotic sieve for primes. *Ann. of Math. (2)*, 148(3):1041–1065, 1998.
- [19] John Friedlander and Henryk Iwaniec. The polynomial  $X^2 + Y^4$  captures its primes. *Ann. of Math. (2)*, 148(3):945–1040, 1998.
- [20] John Friedlander and Henryk Iwaniec. *Opera de cribro*, volume 57 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2010.
- [21] Carl Friedrich Gauss. *Disquisitiones arithmeticae*. Springer-Verlag, New York, 1986. Translated and with a preface by Arthur A. Clarke, Revised by William C. Waterhouse, Cornelius Greither and A. W. Grootendorst and with a preface by Waterhouse.
- [22] Frank Gerth, III. Extension of conjectures of Cohen and Lenstra. *Exposition. Math.*, 5(2):181–184, 1987.
- [23] Franz Halter-Koch, Pierre Kaplan, and Kenneth S. Williams. An Artin character and representations of primes by binary quadratic forms. II. *Manuscripta Math.*, 37(3):357–381, 1982.
- [24] Helmut Hasse. Über die Klassenzahl des Körpers  $P(\sqrt{-2p})$  mit einer Primzahl  $p \neq 2$ . *J. Number Theory*, 1:231–234, 1969.
- [25] Gerald J. Janusz. *Algebraic number fields*. Academic Press [A Subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1973. Pure and Applied Mathematics, Vol. 55.
- [26] Pierre Kaplan. Cycles d’ordre au moins 16 dans le 2-groupe des classes d’idéaux de certains corps quadratiques. *Bull. Soc. Math. France Mém.*, (49-50):113–124, 1977. Utilisation des calculateurs en mathématiques pures (Conf., Limoges, 1975).
- [27] J. C. Lagarias. Signatures of units and congruences (mod 4) in certain real quadratic fields. II. *J. Reine Angew. Math.*, 320:115–126, 1980.

- [28] Jeffrey Clark Lagarias. *The 4-part of the class groups of a quadratic field*. ProQuest LLC, Ann Arbor, MI, 1974. Thesis (Ph.D.)—Massachusetts Institute of Technology.
- [29] Serge Lang. *Algebra*, volume 211 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2002.
- [30] Philip A. Leonard and Kenneth S. Williams. On the divisibility of the class numbers of  $Q(\sqrt{-p})$  and  $Q(\sqrt{-2p})$  by 16. *Canad. Math. Bull.*, 25(2):200–206, 1982.
- [31] D. Milovic. On the 16-rank of class groups of  $\mathbb{Q}(\sqrt{-8p})$  for  $p \equiv -1 \pmod{4}$ . *ArXiv e-prints*, November 2015.
- [32] D. Milovic. The infinitude of  $\mathbb{Q}(\sqrt{-p})$  with class number divisible by 16. *ArXiv e-prints*, February 2015.
- [33] Bernard Oriat. Sur la divisibilité par 8 et 16 des nombres de classes d'idéaux des corps quadratiques  $Q(\sqrt{2p})$  et  $Q(\sqrt{-2})$ . *J. Math. Soc. Japan*, 30(2):279–285, 1978.
- [34] L. Rédei. Arithmetischer Beweis des Satzes über die Anzahl der durch vier teilbaren Invarianten der absoluten Klassengruppe im quadratischen Zahlkörper. *J. Reine Angew. Math.*, 171:55–60, 1934.
- [35] Ladislaus Rédei. Ein neues zahlentheoretisches Symbol mit Anwendungen auf die Theorie der quadratischen Zahlkörper. I. *J. Reine Angew. Math.*, 180:1–43, 1939.
- [36] Hans Reichardt. Zur Struktur der absoluten Idealklassengruppe im quadratischen Zahlkörper. *J. Reine Angew. Math.*, 170:75–82, 1934.
- [37] Lowell Schoenfeld. Sharper Bounds for the Chebyshev Functions  $\theta(x)$  and  $\psi(x)$ . II. *Math. Comp.*, 30(134):337–360, 1976.
- [38] Arnold Scholz. Über die Lösbarkeit der Gleichung  $t^2 - Du^2 = -4$ . *Math. Z.*, 39(1):95–111, 1935.
- [39] Jean-Pierre Serre. Quelques applications du théorème de densité de Chebotarev. *Inst. Hautes Études Sci. Publ. Math.*, (54):323–401, 1981.
- [40] Peter Stevenhagen. Ray class groups and governing fields. In *Théorie des nombres, Année 1988/89, Fasc. 1*, Publ. Math. Fac. Sci. Besançon, page 93. Univ. Franche-Comté, Besançon, 1989.
- [41] Peter Stevenhagen. Divisibility by 2-powers of certain quadratic class numbers. *J. Number Theory*, 43(1):1–19, 1993.
- [42] Peter Stevenhagen. The number of real quadratic fields having units of negative norm. *Experiment. Math.*, 2(2):121–136, 1993.

- [43] Takashi Taniguchi and Frank Thorne. Secondary terms in counting functions for cubic fields. *Duke Math. J.*, 162(13):2451–2508, 2013.
- [44] I. M. Vinogradov. *The method of trigonometrical sums in the theory of numbers*. Dover Publications, Inc., Mineola, NY, 2004. Translated from the Russian, revised and annotated by K. F. Roth and Anne Davenport, Reprint of the 1954 translation.
- [45] Kenneth S. Williams. On the class number of  $\mathbf{Q}(\sqrt{-p})$  modulo 16, for  $p \equiv 1 \pmod{8}$  a prime. *Acta Arith.*, 39(4):381–398, 1981.